

On the X-ray Properties of OH Megamaser Sources: Chandra Snapshot Observations

Cristian Vignali,^{1,2*} William N. Brandt,^{3*} Andrea Comastri^{2*} and Jeremy Darling^{4*}

¹ Dipartimento di Astronomia, Università degli Studi di Bologna, Via Ranzani 1, 40127 Bologna, Italy

² INAF – Osservatorio Astronomico di Bologna, Via Ranzani 1, 40127 Bologna, Italy

³ Department of Astronomy and Astrophysics, The Pennsylvania State University, 525 Davey Laboratory, University Park, PA 16802, USA

⁴ Carnegie Observatories, 813 Santa Barbara Street, Pasadena, CA 91101, USA

Accepted 2005 August 22. Received 2005 July 20

ABSTRACT

We present *Chandra* snapshot observations for a sample of 7 sources selected from the Arecibo OH megamaser (OHM) survey at $z \approx 0.13 - 0.22$ and with far-infrared luminosities in excess of $10^{11} L_{\odot}$. In contrast with the known H_2O megamasers, which are mostly associated with powerful Active Galactic Nuclei (AGN), the situation is far less clear for OHMs, which have been poorly studied in the X-ray band thus far. All of the observed sources are X-ray weak, with only one OHM, IRAS FSC 03521+0028 ($z = 0.15$), being detected by *Chandra* (with 5 counts). The results from this pilot program indicate that the X-ray emission, with luminosities of less than $\approx 10^{42} \text{ erg s}^{-1}$, is consistent with that from star formation (as also suggested in some cases by the optical spectra) and low-luminosity AGN emission. If an AGN is present, its contribution to the broad-band emission of OHM galaxies is likely modest. Under reasonable assumptions about the intrinsic X-ray spectral shape, the observed count distribution from stacking analysis suggests absorption of $\approx 10^{22} \text{ cm}^{-2}$.

Key words: galaxies: active — galaxies: interactions — galaxies: nuclei — galaxies: starburst — X-rays: galaxies

1 INTRODUCTION

Extragalactic hydroxyl (OH) megamaser (OHM) emission has been studied since the early eighties when it was discovered in Arp 220 (Baan, Wood & Haschick 1982). To be produced, OHM activity requires (1) high molecular density ($n_{H_2} = 10^{4-7} \text{ cm}^{-3}$; Baan 1991), (2) a “pump” to invert the hyperfine population of the OH ground state, and (3) a source of 18 cm continuum emission to stimulate maser emission (Burdyuzha & Komberg 1990), whose main lines are at 1667 MHz and 1665 MHz and have luminosities $L_{OH} = 10^{1-4} L_{\odot}$. The galaxy merger environment is able to supply all of these requirements: the merger/interaction concentrates molecular gas in the merger nuclei, creates strong far-infrared (FIR; 8–1000 μm) dust emission from reprocessed starburst and active galactic nucleus (AGN) activity, and produces radio continuum emission. The FIR radiation field and/or collisional shocks in the molecular gas can invert the OH population and thus allow maser emission.

Not surprisingly, all known OHMs have been observed in luminous infrared galaxies (LIRGs; $L_{FIR} > 10^{11} L_{\odot}$), often favouring

the most FIR luminous (e.g., Baan 1991; Baan, Salzer & LeWinter 1998; Darling & Giovanelli 2002a, hereafter DG02a; Lo 2005), the ultraluminous infrared galaxies (ULIRGs; $L_{FIR} > 10^{12} L_{\odot}$). The high FIR luminosities of ULIRGs are commonly thought to arise from dust absorption and FIR re-emission of an intense but obscured starburst and/or AGN radiation field. While it was known early on that ULIRGs nearly always show evidence for interactions (i.e., collisions/mergers; e.g., Sanders & Mirabel 1996; Clements et al. 1996; Borne et al. 2000; Farrah et al. 2001, 2003), many investigators have debated the luminosity dependence of the fraction of interacting systems among the ULIRG population and the possibility that these sources may represent the transition stage between galaxy mergers and quasars (e.g., Farrah et al. 2001; Tacconi et al. 2002; see also Yun et al. 2004).

OHM studies provide powerful diagnostics of the physical conditions in the innermost regions of luminous FIR galaxies. There is evidence that the OH emission is often produced by an ensemble of many masing regions in the nuclear regions of (U)LIRGs on scales of a few hundred parsecs or less (Diamond et al. 1999). VLBI observations of a few nearby OHMs have demonstrated that OH maser emission can arise in circumnuclear discs or tori such as for III Zw 35 (Pihlström et al. 2001) or Mrk 231 (Klöckner, Baan & Garrett 2003), or, as for Arp 220 (Rovilos et al. 2003), the emis-

* E-mail: cristian.vignali@bo.astro.it (CV); niel@astro.psu.edu (WNB); andrea.comastri@bo.astro.it (AC); darling@ociw.edu (JD)

Table 1. *Chandra* Cycles 4 and 5 observations of OH megamaser galaxies: observation log.

Src. Name		Optical		X-ray Obs.	Exp. Time	Reference
IRAS FSC	z	α_{2000}	δ_{2000}	Date	(ks)	(Arecibo Survey)
01562+2528	0.1658	01 59 02.61	+25 42 35.4	2003 Nov. 08–09	4.00	(1)
02524+2046	0.1815	02 55 17.09	+20 58 56.5	2003 Nov. 08	4.31	(1)
03521+0028	0.1522	03 54 42.21	+00 37 03.8	2002 Dec. 25	9.78	(1)
08201+2801	0.1680	08 23 12.62	+27 51 39.6	2003 Nov. 20	4.41	(2)
08279+0956	0.2085	08 30 39.38	+09 46 36.1	2004 Jan. 13	4.38	(2)
09531+1430	0.2151	09 55 50.11	+14 16 07.9	2003 Dec. 09	4.03	(2)
09539+0857	0.1290	09 56 34.30	+08 43 06.1	2004 Jan. 06	4.81	(2)

NOTES — The optical positions have been derived from the Digital Palomar Sky Survey (DPOSS2) *R*-band images using SExtractor (Bertin & Arnouts 1996). REFERENCES — (1) Darling & Giovanelli 2002a (DG02a); (2) Darling & Giovanelli 2001 (DG01).

sion can show a complicated, irregular morphology and velocity structure. Blueshifted/redshifted line components in several OHMs (Baan, Haschick & Henkel 1989; Pihlström et al. 2005) have been interpreted as due to bulk outflows/inflows of molecular gas; this is expected given the tidal streaming and starburst-driven winds associated with major mergers. It is noteworthy that most VLBI studies of OHMs show unresolved maser spots of surprising velocity width (tens of km s^{-1}), often attributed to turbulence and suggestive that compact OHM emission may not be pumped by the FIR radiation field alone (e.g. Lonsdale et al. 1998). A notable exception is the infrared quasar Mrk 231, which lacks a compact maser component contrary to expectation (Lonsdale et al. 2003; Klöckner, Baan & Garrett 2003). The relationship of OHMs to starbursts versus AGN thus remains murky: all manner of masing is seen coupled with all manner of optical classifications. To obtain deeper insight into the OHM phenomenon, we require a minimal-obscuration means to quantify the role of AGN versus starburst in masing regions. The radio-FIR relation hints at some difference between masing and non-masing (U)LIRGs (DG02a), but there is a degeneracy between starbursts and AGN in this relation.

In contrast with H_2O megamasers, which are likely pumped by an AGN X-ray radiation field (e.g., Baan 1997; Braatz, Wilson & Henkel 1997; Henkel et al. 1998, 2005; Townsend et al. 2001; Maloney 2002; Braatz et al. 2003, 2004), the AGN content of OHMs has not been investigated with a minimal-obscuration probe of AGN activity, aside from radio continuum observations of radio-loud AGN that lie well off of the radio-FIR relation. A previous optical study of OH maser sources (most of which were OHMs) found comparable fractions of AGN- and starburst-dominated galaxies (Baan et al. 1998) and a non-negligible fraction of composite spectra (i.e., showing evidence of both AGN and starburst activity) in the OH maser population. Because of their penetrating nature, X-rays represent an efficient tool to provide constraints on the engine powering OHMs. Furthermore, X-rays typically provide maximal contrast between the AGN emission and that of the host galaxy and/or starburst component (e.g., Vignati et al. 1999). Hereafter we adopt $H_0=75 \text{ km s}^{-1} \text{ Mpc}^{-1}$ in a Λ -cosmology ($\Omega_M=0.3$ and $\Omega_\Lambda=0.7$, Spergel et al. 2003).

2 SAMPLE SELECTION AND SOURCE PROPERTIES

The goal of the present paper is to provide basic constraints on the X-ray emission of OHM galaxies using *Chandra* snapshot observations. Past studies have shown evidence for a correlation be-

tween the OH line width and the X-ray luminosity in the soft band (Kandalyan 2003) using a compilation of OH masers from the literature (mostly from Baan et al. 1998) detected by *ROSAT*. This was interpreted as suggesting that X-ray heating of molecular gas provides additional collisional excitation to the maser emission, although further studies are required to investigate this issue. We note, however, that the compilation of OHMs chosen by Kandalyan (2003) is heterogeneous, since it was not drawn from a well defined sample. For these reasons, we prefer to focus, in this pilot *Chandra* program, on the sample of OHMs selected from the Arecibo Observatory OH megamaser survey which is a flux-limited survey designed to quantify the relationships between merging galaxies and the OHMs that they produce, with the goal of using OHMs as luminous tracers of mergers (and, possibly, of star formation; Townsend et al. 2001) at high redshifts (Darling & Giovanelli 2000, 2001, 2002b, hereafter DG00, DG01, and DG02b, respectively). This survey has been conducted over about one-quarter of the sky to a distance limit of roughly 1 Gpc. Candidates for observations with the Arecibo Observatory were selected from the PSCz (Saunders et al. 2000), which is a flux-limited ($IRAS f_{60 \mu\text{m}} > 0.6 \text{ Jy}$) redshift survey of $\approx 15,000$ *IRAS* galaxies over 84 per cent of the sky. The Arecibo survey detected 50 new OHMs in the redshift range 0.11–0.27 (DG02a,b), thus doubling the sample of known OHMs and increasing the $z > 0.1$ sample sevenfold. Thanks to this survey, it is now evident that the fraction of OHMs in (U)LIRGs is an increasing function of L_{FIR} : about one-third of the “warmest” (i.e., with higher dust temperatures) ULIRGs are characterized by OH megamasers (DG02a).

The OHM galaxies proposed for observation with *Chandra* have been chosen using the catalog published by DG01 available at the time of *Chandra* Cycle 5 proposals (6 of the 7 OHMs described in the following); our targets are among the highest luminosity members ($L_{\text{OH}} > 10^3 L_\odot$; ≈ 30 –40 per cent of the OHM population in the Arecibo survey) of the maser population (filled circles in Fig. 1). They also constitute a complete sample, since they were targeted in order of right ascension. One additional OHM, characterized by lower OH luminosity (filled triangle in Fig. 1), has been included using an archival *Chandra* observation. It is worth noting that our OHMs lie at higher redshifts than most of the OHMs with X-ray detections in the literature (e.g., Arp 220 at $z = 0.018$, Mrk 231 at $z = 0.042$, etc.). On the basis of the merger/interaction scenario for OHMs discussed in §1, the present sample, with all of the 7 OHMs being characterized by double nuclei¹ and/or tidal

¹ We note that, given the presence of double nuclei, the optical positions




Figure 1. Logarithm of the OH luminosity vs. FIR luminosity (both in units of $L_{\odot}$) for the Arecibo OHM final sample (DG02b). The filled symbols indicate the OHMs with X-ray observations, while the filled triangle shows the X-ray detected OHM. The short-dashed lines indicate the permitted range of L_{FIR} when the flux density at $100 \mu\text{m}$ is only available as an upper limit (see DG02b); in the case of OHMs with X-ray observations, both circles are filled. Note that all of the OHM galaxies have L_{FIR} in the LIRG regime, with a significant fraction (≈ 35 per cent) being ULIRGs.

tails (Darling et al., in preparation), can be considered representative of the OH megamaser population overall. Furthermore, their IR colour ($\log f_{100}/f_{60}$) distribution shows no significant evidence for any observational bias toward “warm” (i.e., lower f_{100}/f_{60} values, possibly related to a contribution from an active nucleus) or “cold” galaxies (see Fig. 2).

2.1 Notes on individual sources and optical spectra

In the following, we report on the OH line and optical properties of the 7 OHM galaxies with *Chandra* observations. A detailed description of their optical spectra is presented in Darling et al. (in preparation).

01562+2528: this source shows the presence of two nuclei (one much brighter than the other) in the optical (with a separation of $6.3''$, i.e., 17.9 kpc), suggestive of a multiple merger. The Eastern component has a starburst classification, while for the Western component there is not enough signal for a spectral classification (see Table 2 for references).

02524+2046: this object has the most unusual spectrum of the Arecibo OHM survey, showing multiple strong narrow components in both OH lines. The optical morphology is elliptical-like with a

reported in Table 1 are intermediate between the two nuclei; in these cases, both nuclei fall in the same $2''$ -radius aperture used for X-ray manual photometry (see §3). For IRAS FSC 01562+2528, we report the position of the optically most luminous nucleus, given the association of the OH and radio emissions with this nucleus.




Figure 2. Logarithm of the OH luminosity (in units of $L_{\odot}$) vs. IR colour ($\log f_{100}/f_{60}$) for the OHM galaxies from DG02b. Symbols are the same as in Fig. 1.

single tidal tail (Darling et al., in preparation), while the optical spectrum is typical of a starburst galaxy.

03521+0028: this source is comprised of a close pair; both components, separated by $1.3''$ (corresponding to 3.4 kpc), are visible in the Sloan Digital Sky Survey (SDSS) but not in the DPOSS2. This object has been classified as a starburst according to the ISO spectrum (Lutz, Veilleux & Genzel 1999) and a low-ionisation nuclear emission-line region galaxy (LINER) according to the optical spectrum (Nagar et al. 2003).

08201+2801: the *HST* image shows two nuclei (with a separation of $1.1''$, corresponding to 3.2 kpc), having a starburst classification, and a tidal tail, suggesting an advanced merger (DG01). The NVSS flux density reported in DG01 is associated with a source at a distance of $\approx 23''$. Nagar et al. (2003) reported a flux-density upper limit of 0.8 mJy at 15 GHz, roughly corresponding to a 1.4–15 GHz spectral slope of $\gtrsim 0.75$.

08279+0956: the optical image of this source (classified as a LINER) shows the presence of 3 tidal tails (Darling et al., in preparation).

09531+1430: this galaxy is an interacting system whose components are blended in the images from the Automatic Plate Measuring (APM) facility; the 2MASS position is on the Western component (having a Seyfert 2 spectrum), while the FIRST position is on the Eastern component (characterized by a starburst spectrum; Darling et al., in preparation).

09539+0857: the optical image shows the presence of tidal tails and a single nucleus, classified as a LINER (Veilleux, Kim & Sanders 1999) or Seyfert 2 (Darling et al., in preparation). The two classifications were performed in an identical manner. In both cases, the $[\text{O III}]/\text{H}\beta$ ratio lies very close to the (somewhat arbitrary) dividing line between LINERs and Seyfert 2s, and the measured line ratios obtained by both groups agree to within the measurement uncertainties. Nagar et al. (2003) reported a flux density of 1.2 mJy at 15 GHz, corresponding to a 1.4–15 GHz spectral slope of ≈ 0.6 –0.8.

Table 2. Multi-wavelength properties of the Arecibo sample of OHMs observed by *Chandra*.

Source	z	N_{H}	R	B	f_{12}	f_{25}	f_{60}	f_{100}	$\log L_{\text{FIR}}$	$\log L_{\text{OH}}$	$f_{1.4 \text{ GHz}}$	$f_{1.4 \text{ GHz}}$	F_{FB}	$\log L_{2-10}$	Class.	Ref.
			APM		IRAS						FIRST					
(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)	(9)	(10)	(11)	(12)	(13)	(14)	(15)	(16)	(17)
01562+2528	0.1658	7.46	15.58	17.23	< 0.126	< 0.133	0.809	1.623	11.90	3.25		5.9	< 11.3	< 4.4×10^{41}	STB	(1)
02524+2046	0.1815	11.3	18.08	20.02	< 0.055	< 0.079	0.957	< 4.789	11.81–12.28	3.74		2.6	< 11.4	< 5.4×10^{41}	STB	(1)
03521+0028†	0.1522	12.6	18.44	> 21.13	< 0.250	0.233	2.638	3.833	12.28	2.44		6.1	$3.9^{+2.7}_{-1.7}$	1.3×10^{41}	STB/L	(1,2,3,4)
08201+2801†	0.1680	3.59	17.12	18.39	< 0.085	< 0.162	1.171	1.429	12.00	3.45	4.74		< 4.4	< 1.8×10^{41}	STB	(4,5)
08279+0956	0.2085	3.80	18.08	20.04	< 0.079	< 0.192	0.586	< 1.263	11.75–12.01	3.23	2.84	3.7	< 9.5	< 6.1×10^{41}	L	(1)
09531+1430	0.2151	3.07	17.25	> 19.27	< 0.140	< 0.169	< 0.776	1.040	12.08	3.42	3.23	2.5	< 10.2	< 1.7×10^{42}	S2/L††	(1)
09539+0857†	0.1290	3.08	17.74	19.62	< 0.154	< 0.147	1.438	1.044	11.79	3.48	5.32	8.50	< 8.5	< 1.9×10^{41}	S2/L	(1,3)

Luminosities are computed using $H_0=75 \text{ km s}^{-1} \text{ Mpc}^{-1}$, $\Omega_M=0.3$ and $\Omega_\Lambda=0.7$.

† Also in the SDSS Data Release 3 catalog (DR3; Abazajian et al. 2005).

†† The Eastern nucleus is classified as a starburst, while the Western nucleus is classified as a Seyfert 2 (see text for details).

(1) IRAS FSC Name; (2) redshift; (3) Galactic column density from Dickey & Lockman (1990), in units of 10^{20} cm^{-2} ; (4–5) R - and B -band magnitudes from APM; (6–9) IRAS flux densities, in units of Jy; (10) log of the FIR luminosity (in units of $L_\odot$) computed according to $L_{\text{FIR}}=3.96 \times 10^5 \times D_L^2 (2.58 \times f_{60} + f_{100})$, where D_L is the luminosity distance in Mpc (see Fullmer & Lonsdale 1989); when f_{100} is an upper limit, the permitted range of L_{FIR} is reported (see §2.2 of DG02b); (11) log of the OH luminosity, in units of $L_\odot$; (12) integrated 1.4 GHz flux density from FIRST (Becker, White & Helfand 1995); (13) 1.4 GHz flux density from NVSS (Condon et al. 1998); (14) Galactic absorption-corrected flux in the observed 0.5–8 keV band, in units of $10^{-15} \text{ erg cm}^{-2} \text{ s}^{-1}$; (15) 2–10 keV rest-frame luminosity corrected for the effects of Galactic absorption and computed from the full-band flux (or upper limit) assuming $\Gamma = 2.0$, in units of erg s^{-1} ; (16) spectroscopic classification: STB=starburst; L=LINER; S2=Seyfert 2; (17) reference for the spectroscopic classification.

LIST OF REFERENCES — (1) Darling et al., in preparation; (2) Lutz, Veilleux & Genzel 1999; (3) Veilleux, Kim & Sanders 1999; (4) Nagar et al. 2003; (5) Kim, Veilleux & Sanders 1998.

3 X-RAY OBSERVATIONS AND DATA REDUCTION

Six out of the 7 OHMs presented here were targeted by *Chandra* during Cycle 5. IRAS FSC 03521+0028, found at the aim point of a *Chandra* archival observation, was observed in Cycle 4. This source was targeted in a *Chandra* program aimed at observing a sample of ULIRGs (see Teng et al. 2005); we are confident that the inclusion of this object does not bias our “original” sample in terms of AGN content (see §4). The observation log is shown in Table 1.

All of the sources were observed with the Advanced CCD Imaging Spectrometer (ACIS; Garmire et al. 2003) with the S3 CCD at the aimpoint. Standard data reduction was adopted (see §2 of Vignali et al. 2005 for a detailed description) using the *Chandra* Interactive Analysis of Observations (CIAO) Version 3.2 software. Source detection was carried out with WAVDETECT (Freeman et al. 2002) using a false-positive probability threshold of 10^{-6} . We searched for *Chandra* sources within $\approx 0.8''$ of the optical position. IRAS FSC 03521+0028 is the only detected source (filled triangle in Fig. 1), with 5 counts in the observed 0.5–8 keV band; we note that this is also the source with the highest exposure by *Chandra* (see Table 1). Given the adopted threshold and the small number of pixels being searched due to the known source positions and the sub-arcsec on-axis angular resolution of *Chandra*, the probability that this detection is spurious is extremely low ($\approx 8 \times 10^{-6}$). All of the remaining sources (shown as filled circles in Fig. 1) were not detected also using higher (i.e., less conservative) threshold values (up to 10^{-4}).

Table 3 summarizes the X-ray photometric results in the soft band (0.5–2 keV), the hard band (2–8 keV), and the full band (0.5–8 keV) using circular apertures centred on the optical source position and an extraction radius of $2''$. Although two X-ray photons falling in adjacent pixels can be considered in some circumstances a detection (given the low background level of *Chandra* observations; see, e.g., Vignali et al. 2005), we prefer to follow a more conservative approach and report these sources (2 in the present sample; see Table 3) as X-ray undetected.

The final catalog of OHMs from the Arecibo survey (DG02b) was also cross correlated with the *ROSAT* and *XMM-Newton* archives; we obtained no further X-ray detections. X-ray constraints from *ROSAT* observations (available for 9 OHMs) are generally loose because of the large off-axis angles of the sources

Table 3. X-ray counts and full-band count rates.

Source	X-ray Counts			CR
	(0.5–2 keV)	(2–8 keV)	(0.5–8 keV)	(0.5–8 keV)
01562+2528	< 3.0	< 6.4	< 6.4★	< 1.60
02524+2046	< 4.8	< 4.8	< 6.4	< 1.49
03521+0028	$3.0^{+2.9}_{-1.6}$	< 6.4	$4.9^{+3.4}_{-2.1}$	$0.50^{+0.35}_{-0.22}$
08201+2801	< 3.0	< 3.0	< 3.0	< 0.68
08279+0956	< 6.4	< 3.0	< 6.4★	< 1.46
09531+1430	< 3.0	< 6.4	< 6.4	< 1.59
09539+0857	< 6.4	< 3.0	< 6.4	< 1.33

★ The two X-ray counts are contiguous.

Errors on the X-ray counts were computed according to Gehrels (1986). The upper limits are at the 95% confidence level and were computed according to Kraft, Burrows & Nousek (1991). For the sake of clarity, upper limits of 3.0, 4.8, and 6.4 indicate that 0, 1, and 2 X-ray counts, respectively, have been found within an extraction region of radius $2''$ centred on the position of the OHM source (considering the background within this source extraction region to be negligible). Count rates (CR) are in units of $10^{-3} \text{ counts s}^{-1}$.

and the shallow exposures of the pointed PSPC and HRI observations; hence these constraints will not be used in this paper. One OHM was targeted by *XMM-Newton* (IRAS FSC 13218+0552 at $z = 0.205$, with net exposures of ≈ 8.6 – 9.7 ks in the EPIC MOS/pn cameras) but only an upper limit, consistent with the values we obtained for our sources (see §4 and Table 2), was derived (Bianchi et al. 2005). Recently, it has been suggested that this source, characterized by an extremely red infrared continuum (Low et al. 1989) and a blueshifted component in the [O III] line (e.g., Zheng et al. 2002), might harbour a Compton-thick AGN (i.e., with $N_{\text{H}} \gtrsim 10^{24} \text{ cm}^{-2}$), surrounded by a strong starburst (see §4.4 and Fig. 6 of Bianchi et al. 2005). We also note that IRAS FSC 13218+0552 has the warmest f_{100}/f_{60} colour in Fig. 2 (i.e., it is the left-most object in the plot), suggestive of an AGN contribution at IR wavelengths.




Figure 3. Rest-frame FIR luminosity vs. X-ray luminosity for the OHM galaxies presented in this paper (filled circles: X-ray upper limits; filled triangle: X-ray detected OHM; filled square: stacked X-ray counts from the whole sample of 7 OHMs observed by *Chandra*, see §4.1). The open triangle indicates the OHM from the Arecibo sample which was observed but not detected by XMM-Newton (see Bianchi et al. 2005 and §3). For comparison, some other sources are reported (after correcting, when possible, for intrinsic absorption in the X-ray band): the sub-mm galaxies in the *Chandra* Deep Field-North (Alexander et al. 2003, 2005a,b) likely containing an AGN (open circles) and a “pure” starburst (small open squares); well known AGN-dominated (mostly in the right-hand side of the plot) and star-formation dominated sources (in the left-most region of the figure) from the literature plus two distant well-studied sub-mm galaxies (SMM 02399–0134 and SMM 02399–0136), all shown as large open squares with labels. The slanted lines show ratios of constant X-ray to FIR luminosity; this ratio is typically ≈ 3 –30 per cent for AGN-dominated sources. Adapted from Alexander et al. (2003); see Alexander et al. (2005a) for details. Note that some of the sources from the literature sample are low-redshift OHM galaxies (marked by a “*” after the source name).

4 WHAT ARE X-RAYS TELLING US ABOUT THE NATURE OF THE OH MEGAMASERS?

The broad-band properties of the OHM sample targeted by *Chandra* are reported in Table 2. Both the photon index and X-ray luminosity can provide constraints on the nature of the high-energy emission from OHM galaxies. However, such constraints, with the current data, are basic due to the low counting statistics. All of the X-ray sources have rest-frame 2–10 keV luminosities at most of $\approx 10^{42}$ erg s $^{-1}$ (see Table 2), which is consistent with both star formation (e.g., Zezas, Alonso-Herrero & Ward 2001; Alexander et al. 2002) and low-luminosity AGN (LLAGN) activity (e.g., Terashima et al. 2002; Terashima & Wilson 2003). The possibility that the observed X-ray emission is largely due to star formation finds support from Fig. 3, where the FIR luminosity is plotted vs. the 0.5–8 keV luminosity. Our sources are located in the left-most part of the diagram, where either starbursts (e.g., Arp 220) or starburst-dominated (i.e., with little apparent contribution from the AGN to the bolometric emission; see Alexander et al. 2005a) sub-mm sources typically lie. However, some X-ray obscured AGN whose broad-band emission is dominated by star formation are also present in this re-

gion; a fraction of these sources are lower redshift OHM sources (e.g., Mrk 273 and UGC 5101); the possibility that our sources could contain absorbed AGN will be discussed in detail using X-ray stacking analysis (see §4.1).

Optical/near-infrared (NIR) spectroscopy (see §2.1) indicates that 50 per cent of our sources show the typical signatures of starburst galaxies, 25 per cent have a LINER classification, and 25 per cent a Seyfert 2 classification (according to the classification reported in Darling et al., in preparation; see also column (16) in Table 2); IRAS FSC 09531+1430 is counted twice because both nuclei were spectroscopically identified (see Darling et al., in preparation). We note, however, that the optical classification of these sources is difficult: when more than one nucleus is present, unless the projected physical separation of the nuclei is larger than the resolution of the spectrograph, optical spectra will be a blend of nuclei, dominated by the optically dominant (i.e., the least obscured and/or most optically luminous in emission lines) nucleus. However, OHMs seem to favour the dustiest environments and may select the optically “subordinate” nucleus. Therefore, there may be little correlation between OHM properties and optical spectral type. Furthermore, we note that optical and NIR studies are sometimes not adequate to unveil the engine of the X-ray emission. An emblematic, though perhaps extreme, case of different classification at different wavelengths is provided by the ULIRG NGC 6240, which is classified as a starburst in the NIR (Genzel et al. 1998), a LINER in the optical (Veilleux et al. 1995) and a Compton-thick AGN in the hard X-ray band (Vignati et al. 1999; also see Komossa et al. 2003). Some further cases similar to that of NGC 6240 are presented by Maiolino et al. (2003).

Finally, it is interesting to note that the X-ray luminosity and the X-ray-to-FIR luminosity ratio of the only X-ray detected OHM in the present sample do not appear to be different from those of the other sources; given the number of observed counts (see Table 3), the detection of this source has been possible because of the exposure time which is double that of the other OHMs.

4.1 X-ray stacking analysis

We can place tighter constraints on the average X-ray properties of the OHMs under investigation by “stacking” their counts. Our sources have $8^{+4.0}_{-2.8}$ counts in the 0.5–2 keV band and $7^{+3.8}_{-2.6}$ counts in the 2–8 keV band. Monte Carlo simulations in regions close to our targets (see Vignali et al. 2005 for details about the adopted procedure) indicate that the background in all of the 7 fields does not contribute more than 0.14 and 0.31 counts in the soft and hard bands, respectively; therefore we are confident we obtain secure detections in both bands by stacking the counts from the 7 OHMs.² The ratio of the observed hard vs. soft-band source counts ($0.85^{+0.65}_{-0.43}$) suggests that our sources have, on average, a photon index flatter ($\approx 0.7^{+0.6}_{-0.5}$) than $\Gamma=1.8$ –2.1, which is typically used for AGN and starburst galaxies at hard X-ray energies (e.g., Piconcelli et al. 2005; Ptak et al. 1999).³ If the hardness of our sources is due to the presence of intrinsic absorption, the implied average column density is $\approx 1.5 \times 10^{22}$ cm $^{-2}$ (ranging between $\approx 5.8 \times 10^{21}$ cm $^{-2}$

² The Poisson probability of obtaining 8 (7) counts or more when 0.14 (0.31) counts are expected is $\approx 3.2 \times 10^{-12}$ ($\approx 4.2 \times 10^{-8}$).

³ We note, however, that a large population of neutron star high-mass X-ray binaries, whose presence is expected in cases of galaxies undergoing intense star formation, can produce a flat ($\Gamma \approx 0.5$ –1) photon index; see Colbert et al. (2004).

and $\approx 2.6 \times 10^{22} \text{ cm}^{-2}$) at the average redshift of $z = 0.17$. The corresponding absorption-corrected rest-frame 2–10 keV luminosity would be in the range $\approx 1\text{--}3 \times 10^{41} \text{ erg s}^{-1}$. Intrinsically flatter X-ray slopes would produce lower column densities (by ≈ 20 per cent, if $\Gamma = 1.6$ instead of 2.0 is adopted), while the presence of an additional soft X-ray component (e.g., thermal emission from the starburst) would produce the opposite effect.

Figure 3 shows that, if an active nucleus is present in the OHMs under investigation, its contribution to their broad-band emission is probably modest, unless the AGN is Compton-thick. In the Compton-thick case, the X-ray emission in the *Chandra* band would be the small (presumably a few per cent; see, e.g., Comastri 2004; Guainazzi, Matt & Perola 2005) scattered/reflected fraction of the nuclear X-ray emission, which would be $\lesssim$ a few $\times 10^{43} \text{ erg s}^{-1}$ (i.e., at most in the Seyfert regime). The reprocessing of the X-ray and ultraviolet emissions at longer wavelengths would still be insufficient to produce the observed FIR luminosity without requiring a dominant contribution from star formation. We note that some sources from the literature sample are characterized by Compton-thick absorption (e.g., Mrk 231, Gallagher et al. 2002, 2005, Braitto et al. 2004; UGC 5101, Maiolino et al. 2003).

Although the presence of obscuration is uncertain and needs to be confirmed by deeper observations with *Chandra* for an extended sample of OHMs and, possibly, with *XMM-Newton* for the most “promising” sources, the column density allowed by the current data is consistent with the X-ray absorption observed in many H_2O megamasers. These sources, however, are mostly associated with X-ray bright ($\approx 10^{42\text{--}43} \text{ erg s}^{-1}$) and obscured ($N_{\text{H}} \gtrsim 10^{22} \text{ cm}^{-2}$) Seyfert 2 galaxies (e.g., Maloney 2002; Braatz et al. 2003), where the amplifying water vapor molecules are most likely pumped by collisional processes resulting from X-ray heating in parsec-scale molecular tori with densities of $\approx 10^{7\text{--}8} \text{ cm}^{-3}$ (e.g., Neufeld, Maloney & Conger 1994). Column densities of $\approx 10^{22} \text{ cm}^{-2}$ are not uncommon in local (U)LIRGs (e.g., Risaliti et al. 2000) and LLAGN (e.g., Terashima et al. 2002; Terashima & Wilson 2003). Given the presence of double nuclei and/or indications of past mergers in the OHMs of our sample, it is possible that the starburst itself can produce the obscuration, as suggested by some authors (e.g., Fabian et al. 1998; Ohsuga & Unemura 2001; Wada & Norman 2002).

4.2 X-ray observations of “bona-fide” OHMs from the literature

We obtained further constraints on the X-ray emission from OHM galaxies using the sample of 54 “bona-fide” OH masers [both kilo- ($L_{\text{OH}} \leq 10^1 L_{\odot}$) and mega-] from the literature (mostly from Baan et al. 1998), listed in Tables 7 and 8 of DG02a. We caution the reader against over-interpretation of the results we present in this section: this sample is not well defined, lacks either published OH spectra or measurements of the 1667 MHz OH line in about 50 per cent of the cases, and includes some suspect OH detections, as pointed out by DG02a.

Twenty-four OH masers have accessible pointed X-ray observations in the *ROSAT*, *Chandra*, and *XMM-Newton* archives (for one additional source, the *XMM-Newton* data are not public yet); all of these galaxies are OHMs. Seventeen sources are X-ray detected; the high detection rate (≈ 70 per cent) is due to the fact that most of these sources were targeted by the sensitive X-ray instruments on-board *Chandra* and *XMM-Newton*, while the X-ray non-detections are mostly due to *ROSAT* observations where the sources are located at large off-axis angles. For most of the sources with

sensitive observations, X-ray information is available from published work (Gallagher et al. 2002, 2005; Xia et al. 2002; Maiolino et al. 2003; Ptak et al. 2003; Franceschini et al. 2003, hereafter F03; Satyapal, Sambruna & Dudik 2004; Ballo et al. 2004; Braitto et al. 2004; Imanishi & Terashima 2004; Iwasawa et al. 2005); for a couple of sources, *Chandra* data have been retrieved from the archive and analysed using standard procedures.

We find that AGN emission (with intrinsic X-ray luminosities above $\approx 10^{42} \text{ erg s}^{-1}$) is present in eight OHMs. Four of these OHMs are likely characterized by Compton-thick absorption (UGC 05101, Maiolino et al. 2003; NGC 3690 in the galaxy pair Arp 299 system, Della Ceca et al. 2002, Ballo et al. 2004; NGC 4418, Maiolino et al. 2003; Mrk 231, Gallagher et al. 2002, 2005; Braitto et al. 2004). However, Compton-thick absorption cannot be ruled out in some low signal-to-noise ratio sources and in Arp 220 (Iwasawa et al. 2001, 2005).

To provide a more appropriate comparison with our *Chandra* targets, from the subsample of literature OHM with X-ray coverage, we selected those within the same redshift interval, the same L_{OH} range, or the same L_{FIR} range as our seven OHMs observed by *Chandra*. The paucity of objects in the same redshift range (two, with only one X-ray detection) does not allow for any reasonable comparison with our sample; however, it highlights the relevance of the present work, where the redshift range of OHMs with X-ray constraints has been extended significantly. If we select the literature OHMs using the same interval of OH or FIR luminosities as our targets, we find a larger number of objects (12 and nine, respectively). AGN emission is clearly present in three objects (one Compton-thick, Mrk 231, and the other two Compton-thin, Mrk 273 and PKS B1345+145), corresponding to $\approx 25\text{--}33$ per cent of the chosen subsamples. LLAGN could still be present in some of the remaining sources, although the X-ray emission from these objects can be well explained by a thermal gaseous component from a starburst plus a hard, moderately absorbed power-law component due to the unresolved binary population (e.g., F03; Teng et al. 2005).

As a further test, we compared the X-ray spectral results for our seven OHMs (see §4.1) with those obtained by F03 with *XMM-Newton* for their five “bona-fide” lower redshift OHMs without a dominant AGN component (i.e., excluding Mrk 231). We found that our stacked spectrum is generally harder than the X-ray spectra of the F03 OHMs. This result can be ascribed to the different spatial resolutions of the *Chandra* ACIS and *XMM-Newton* EPIC cameras and different source-extraction regions. To test this possibility, we used a sample of three starburst galaxies observed by *Chandra* (Ptak et al. 2003) and *XMM-Newton* (F03). The soft thermal component appears spatially more extended than the hard component, i.e., it likely provides a stronger contribution to the broad-band X-ray spectrum if large source-extraction regions are chosen (as in the case of *XMM-Newton* spectra, extracted from circular regions of radius $20''$). Motivated by these considerations, we rescaled the source-extraction regions of our OHMs to match those adopted by F03 (taking into account the different redshift range of the two samples), obtaining a somewhat softer X-ray spectrum ($\Gamma \approx 1.6 \pm 0.3$), consistent with F03 results.

In the light of the results obtained from the study of the literature OHMs and keeping in mind all the caveats described above, starburst emission with moderate absorption probably within the binary population is likely to explain the average X-ray properties of our seven OHMs (§4.1).

SUMMARY

We have analysed *Chandra* snapshot observations for a sample of 7 galaxies selected from the Arecibo OHM survey at $z \approx 0.13 - 0.22$ and with FIR luminosities in excess of $10^{11} L_{\odot}$. The principal results of this work are

- All of the observed sources are X-ray weak; only one OHM, IRAS FSC 03521+0028 ($z = 0.15$), was detected by *Chandra*, with 5 counts in the 0.5–8 keV band.
- The X-ray emission is consistent with that from star formation (as also suggested by the comparison with most of the literature OHMs with X-ray constraints) and LLAGN; however, if an AGN is present, its contribution to the broad-band emission of OHM galaxies is probably modest.
- Under reasonable assumptions about the intrinsic X-ray spectral shape, the observed count distribution from stacking analysis suggests absorption of $\approx 10^{22} \text{ cm}^{-2}$.

ACKNOWLEDGMENTS

We acknowledge financial support from MIUR (COFIN grant 03-02-23; CV and AC), *Chandra* X-ray Center grant G04-5104X (CV and WNB), and NASA LTSA grant NAG5-13035 (WNB). The authors thank D. Alexander for providing us with the data shown in Fig. 3, S. Bianchi and G. Matt for sharing information on the OHM observed by XMM-Newton before publication, C. Gruppi for useful discussions, A. Tarchi for a careful reading of the manuscript, and the referee for his/her useful comments.

REFERENCES

- Abazajian K. et al., 2005, *AJ*, 129, 1755
- Alexander D.M., Aussel H., Bauer F.E., Brandt W.N., Hornschemeier A.E., Vignali C., Garmire G.P., Schneider D.P., 2002, *ApJ*, 568, L85
- Alexander D.M. et al., 2003, *AJ*, 125, 383
- Alexander D.M., Bauer F.E., Chapman S.C., Smail I., Blain A.W., Brandt W.N., Ivison R.J., 2005a, *ApJ*, in press (astro-ph/0506608)
- Alexander D.M., Smail I., Bauer F.E., Chapman S.C., Blain A.W., Brandt W.N., Ivison R.J., 2005b, *Nature*, 434, 738
- Baan W.A., Wood P.A.D., Haschick A.D., 1982, *ApJ*, 260, L49
- Baan W.A., Haschick A.D., Henkel C., 1989, *ApJ*, 346, 680
- Baan W.A., 1991, in Haschick A., Ho P., eds, *ASP Conf. Ser. Vol. 16, Atoms, Ions, and Molecules: New Results in Spectral Line Astrophysics*, Astron. Soc. Pac., San Francisco, p. 45
- Baan W.A., 1997, *Nature*, 386, 445
- Baan W.A., Salzer J.J., LeWinter R.D., 1998, *ApJ*, 509, 645
- Ballo L., Braitto V., Della Ceca R., Maraschi L., Tavecchio F., Dadina M., 2004, *ApJ*, 600, 634
- Becker R.H., White R.L., Helfand D.J., 1995, *ApJ*, 450, 559
- Bertin E., Arnouts S., 1996, *A&AS*, 117, 393
- Bianchi S., Guainazzi M., Matt G., Chiaberge M., Iwasawa K., Fiore F., Maiolino R., 2005, *MNRAS*, in press (astro-ph/0507323)
- Borne K.D., Bushouse H., Lucas R.A., Colina L., 2000, *ApJ*, 529, L77
- Braatz J.A., Wilson A.S., Henkel C., 1997, *ApJS*, 110, 321
- Braatz J.A., Wilson A.S., Henkel C., Gough R., Sinclair M., 2003, *ApJS*, 146, 249
- Braatz J.A., Henkel C., Greenhill L.J., Moran J.M., Wilson A.S., 2004, *ApJ*, 617, L29
- Braitto V. et al., 2004, *A&A*, 420, 79
- Burdyuzha V.V., Komberg B.V., 1990, *A&A*, 234, 40
- Clements D.L., Sutherland W.J., McMahon R.G., Saunders W., 1996, *MNRAS*, 279, 477
- Colbert E.J.M., Heckman T.M., Ptak A.F., Strickland D.K., Weaver K.A., 2004, *ApJ*, 602, 231
- Comastri A., 2004, in Barger A.J., eds, *Supermassive Black Holes in the Distant Universe*. Kluwer Academic Publishers, Dordrecht, The Netherlands, p. 245
- Condon J.J., Cotton W.D., Greisen E.W., Yin Q.F., Perley R.A., Taylor G.B., Broderick J.J., 1998, *AJ*, 115, 1693
- Darling J., Giovanelli R., 2000, *AJ*, 119, 3003 (DG00)
- Darling J., Giovanelli R., 2001, *AJ*, 121, 1278 (DG01)
- Darling J., Giovanelli R., 2002a, *AJ*, 124, 100 (DG02a)
- Darling J., Giovanelli R., 2002b, *ApJ*, 572, 810 (DG02b)
- Darling J. et al., in preparation
- Della Ceca R. et al. 2002, *ApJ*, 581, L9
- Diamond P.J., Lonsdale C.J., Lonsdale C.J., Smith H.E., 1999, *ApJ*, 511, 178
- Dickey J.M., Lockman F.J., 1990, *ARA&A*, 28, 215
- Fabian A.C., Barcons X., Almaini O., Iwasawa K., 1998, *MNRAS*, 297, L11
- Farrah D. et al., 2001, *MNRAS*, 326, 1333
- Farrah D., Afonso J., Efstathiou A., Rowan-Robinson M., Fox M., Clements D., 2003, *MNRAS*, 343, 585
- Franceschini A. et al., 2003, *MNRAS*, 343, 1181 (F03)
- Freeman P.E., Kashyap V., Rosner R., Lamb D.Q., 2002, *ApJS*, 138, 185
- Fullmer L., Lonsdale C.J., 1989, *Catalogued Galaxies and Quasars Observed in the IRAS Survey, Version 2*, Pasadena (JPL)
- Gallagher S.C., Brandt W.N., Chartas G., Garmire G.P., Sambruna R.M., 2002, *ApJ*, 569, 655
- Gallagher S.C., Schmidt G.D., Smith P.S., Brandt W.N., Chartas G., Hylton S., Hines D.C., Brotherton M.S., 2005, *ApJ*, in press (astro-ph/0506616)
- Garmire G.P., Bautz M.W., Ford P.G., Nousek J.A., Ricker G.R., 2003, *Proc. SPIE*, 4851, 28
- Gehrels N., 1986, *ApJ*, 303, 336
- Genzel R. et al., 1998, *ApJ*, 498, 579
- Guainazzi M., Matt G., Perola G.C., 2005, *A&A*, in press (astro-ph/0508265)
- Henkel C., Wang Y.P., Falcke H., Wilson A.S., Braatz J.A., 1998, *A&A*, 335, 463
- Henkel C., Peck A.B., Tarchi A., Nagar N.M., Braatz J.A., Castangia P., Moscadelli L., 2005, *A&A*, 436, 75
- Kandalyan R.A., 2003, *A&A*, 404, 513
- Kim D.-C., Veilleux S., Sanders D.B., 1998, *ApJ*, 508, 627
- Klößner H.-R., Baan W.A., Garrett M.A., 2003, *Nature*, 421, 821
- Klößner H.-R., Baan W.A., 2004, *A&A*, 419, 887
- Komossa S., Burwitz V., Hasinger G., Predehl P., Kaastra J.S., Icke Y., 2003, *ApJ*, 582, L15
- Kraft R.P., Burrows D.N., Nousek J.A., 1991, *ApJ*, 374, 344
- Imanishi M., Terashima Y., 2004, *AJ*, 127, 758
- Iwasawa K., Matt G., Guainazzi M., Fabian A.C., 2001, *MNRAS*, 326, 894
- Iwasawa K., Sanders D.B., Evans A.S., Trentham N., Miniutti G., Spoon H.W.W., 2005, *MNRAS*, 357, 565
- Lo K.Y., 2005, *ARA&A*, 43, in press
- Lonsdale C.J., Diamond P.J., Smith H.E., Lonsdale C.J., 1998, *ApJ*, 493, L13
- Lonsdale C.J., Lonsdale C.J., Smith H.E., Diamond P. J., 2003, *ApJ*, 592, 804
- Low F.J., Cutri R.M., Kleinmann S.G., Huchra J.P., 1989, *ApJ*, 340, L1
- Lutz D., Veilleux S., Genzel R., 1999, *ApJ*, 517, L13
- Maiolino R. et al., 2003, *MNRAS*, 344, L59
- Maloney P.R., 2002, *PASA*, 19, 401
- Nagar N.M., Wilson A.S., Falcke H., Veilleux S., Maiolino R., 2003, *A&A*, 409, 115
- Neufeld D.A., Maloney P.R., Conger S., 1994, *ApJ*, 436, L127
- Ohsuga K., Umemura M., 2001, *ApJ*, 559, 157
- Piconcelli E., Jimenez-Bailón E., Guainazzi M., Schartel N., Rodríguez-Pascual P.M., Santos-Lleó M., 2005, *A&A*, 432, 15
- Pihlström Y.M., Conway J.E., Booth R.S., Diamond P.J., Polatidis A.G., 2001, *A&A*, 377, 413
- Pihlström Y.M., Baan W.A., Darling J., Klößner H.-R., *ApJ*, 618, 705
- 2005, *ApJ*, 618, 705

Ptak A., Serlemitsos P., Yaqoob T., Mushotzky R., 1999, ApJS, 120, 179

Ptak A., Heckman T., Levenson N.A., Weaver K., Strickland D., 2003, ApJ, 592, 782

Risaliti G., Gilli R., Maiolino R., Salvati M., 2000, A&A, 357, 13

Rovilos E., Diamond P.J., Lonsdale C.J., Lonsdale C.J., Smith H.E., 2003, MNRAS, 342, 373

Sanders D.B., Mirabel I.F., 1996, ARA&A, 34, 749

Satyapal S., Sambruna R.M., Dudik R.P., 2004, A&A, 414, 825

Saunders W. et al., 2000, MNRAS, 317, 55

Spergel D.N. et al., 2003, ApJS, 148, 175

Tacconi L.J., Genzel R., Lutz D., Rigopoulou D., Baker A.J., Iserlohe C., Tecza M., 2002, ApJ, 580, 73

Teng S.H., Wilson A.S., Veilleux S., Young A.J., Sanders D.B., Nagar N.M., 2005, ApJ, in press (astro-ph/0508112)

Terashima Y., Iyomoto N., Ho L.C., Ptak A.F., 2002, ApJS, 139, 1

Terashima Y., Wilson A.S., 2003, ApJ, 583, 145

Townsend R.H.D., Ivison R.J., Smail I., Blain A.W., Frayer D.T., 2001, MNRAS, 328, L17

Veilleux S., Kim D.-C., Sanders D.B., Mazzarella J.M., Soifer B.T., 1995, ApJS, 98, 171

Veilleux S., Kim D.-C., Sanders D.B., 1999, ApJ, 522, 113

Vignali C., Brandt W.N., Schneider D.P., Kaspi S., 2005, AJ, 129, 2519

Vignati P. et al., 1999, A&A, 349, L57

Wada K., Norman C.A., 2002, ApJ, 566, L21

Xia X.Y., Xue S.J., Mao S., Boller T., Deng Z.G., Wu H., 2002, ApJ, 564, 196

Yun M.S., Reddy N.A., Scoville N.Z., Frayer D.T., Robson E.I., Tilanus R.P.J., 2004, ApJ, 601, 723

Zezas A., Alonso-Herrero A., Ward M.J., 2001, Ap&SS, 276, 601

Zheng X.Z., Xia X.Y., Mao S., Wu H., Deng Z.G., 2002, AJ, 124, 18